

The Buzzards Bay National Estuary Program
Pocket Guide to Hydric
Soils for Wetland
Delineations in
Massachusetts



The Buzzards Bay National Estuary Program
September, 2013

This document is largely taken from Delineating Bordering Vegetated Wetlands under the Wetlands Protection Act, published by the Massachusetts Department of Environmental Protection, Division of Wetlands and Waterways and the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0), published by the U.S. Army Engineer Research and Development Center.

This document also includes information from the Field Indicators of Hydric Soils in the United States, A Guide for Identifying and Delineation Hydric Soils, Version 7.0, including the 2013 Errata, as well as Field Indicators for Identifying Hydric Soils in New England, 3rd ed.

This document is meant to be a companion to “The Buzzards Bay National Estuary Program Pocket Guide to Delineating Wetlands,” available at the Buzzards Bay National Estuary Program website, www.buzzardsbay.org.

Unless otherwise noted, photos are from the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0).

Table of Contents

Preface	i
DEP Short List	1
1. Histosols	2
2. Histic Epipedon	3
3. Sulfidic Materials	4
4. Gleyed Matrix	5
5. Depleted Matrix (1)	7
6. Depleted Matrix (2)	8
7. Redox Depletions	9
DEP Difficult Soils	10
1. Sandy Soils	11
2. Floodplain Soils	12
3. Soil from Highly Colored Parent Material	13
4. A-Horizons that are Thick and Very Dark	15
5. Evergreen Forest Soils	16
6. Areas Where the Hydrology has been Recently Altered	18
DEP Particularly Difficult Soils	19
A1 Histosol	21
A5 Stratified Layers	22
A11 Depleted Below Dark Surface	24
A12 Thick Dark Surface	26
S1 Sandy Mucky Mineral	27
S6 Stripped Matrix	28
S7 Dark Surface	29
S8 Polyvalue below Surface	30
S9 Thin Dark Surface	31
F3 Depleted Matrix	32
F6 Redox Dark Surface	34
F7 Depleted Dark Surface	35

DEP Particularly Difficult Soils – cont.

X. Sandy with Redox & Thick, Very Dark Surface	36
XI. B. Redox Depletions & Thick Very Dark Surface	37
F8 Redox Depressions	38
A10 2cm Muck	39
A16 Coastal Prairie Redox	40
S3 Mucky Peat or Peat	41
F12 Iron-Manganese Masses	42
F19 Piedmont Floodplain Soils	43
F21 Red Parent Material.(1)	44
TF2 Red Parent Material (2)	46
TF12 Very Shallow Dark Surface	47
TA6 Mesic Spodic	48
IX A. Spodosols	49
IX B. Spodosols	51
IX C. Spodosols	52
VII. Dark Surface Materials	54
XI. A. Redox Depletions	56
XII. Chroma 3 with Very Dark Surface	57
Appendix A - MLRA Boundaries	58
Appendix B - Thickness Criteria For Particularly Difficult Soils	59
Appendix C - Spodosol Field Criteria	60
Appendix D - Mucky Mineral Field Identification For Particularly Difficult Soils	62
Appendix E - Redox Features- Faint vs. Distinct	63
Appendix F - ACOE Memo regarding use of “Field Indicators for Identifying Hydric Soils in New England, Version 3”	64
Appendix G – Observing and Recording Hue, Value and Chroma	65
Postscript: Indicators of Hydrology	65

Preface

Since 1995, the use of Hydric Soils has been an important part of the delineation process in wetland delineation pursuant to the MGL Chapter 131, section 40, the Massachusetts Wetlands Protection Act.

Hydric Soils are considered a confirmation of wetland hydrology as called for in the Regulatory definition of a bordering vegetated wetland found in 310 CMR 10.55 (2)(c). and should be part of every wetland delineation.

In disturbed sites, the presence of hydric soils is sufficient to determine that “there are indicators of saturated or inundated conditions sufficient to support a predominance of wetland indicator plants.”

For areas that have been recently drained, DEP has determined that “hydric soils are often the best indicators for delineating recently drained wetlands.”

Areas where vegetation has been altered or removed - such as golf courses, lawns, and agricultural fields - require the use of soils and other indicators of hydrology to delineate BVW boundaries. In some cases, such as where vegetation has been cut or removed (e.g. ongoing forestry activity), remnant vegetation should be considered, but other indicators of hydrology also should be used to establish the BVW boundary.

Areas where fill has been placed in wetlands require the analysis of soils directly beneath the fill.

The DEP Wetland Delineation Manual, Delineating Bordering Vegetated Wetlands under the Wetlands Protection Act, published by the Massachusetts Department of Environmental Protection, Division of Wetlands and Waterways, has a list of “some hydric soil indicators” on page 29. These seven indicators have been referenced in this guide as the “DEP Short List.”

On pages 30 and 31 of the Manual, DEP lists six soil types that are hard to analyze. Most of these soils have suggested hydric morphologies and are listed in this guide as “DEP Difficult Soils.”

In the first paragraph on page 30, DEP states, “In particularly difficult cases, consultation with the Natural Resources Conservation Service is recommended.” NRCS uses the latest federal hydric soil list. A list of these soils can be found in

the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0), which is based on Field Indicators of Hydric Soils of the United States, Version 7¹. These soils, if they differ from the “DEP Short List” or the “DEP Difficult Soils” have been listed under the heading “DEP Particularly Difficult Soils.”

Those soils in the Regional Supplement that are already included in the “DEP Short List” are not listed the DEP Particularly Difficult Soils” section. This includes Indicators A2, - Histic Epipedon, A3 - Black Histic, S4 - Sandy Gleyed Matrix, S5 - Sandy Redox, and F2 - Loamy Gleyed Matrix.

Delineations by their nature take place at the transition from hydric to non-hydric soils. So it should not be considered unusual for the delineator to encounter problem areas. In addition to the hydric soil morphologies found in the DEP Manual and the Regional Supplement, practitioners should be aware that the use of Field Indicators for Identifying Hydric Soils in New England, Version 3 is valid at problem sites. This is further explained in a March 11, 2011 memo from Ruth M. Ladd, Chief of the Policy Analysis and Technical Support Branch of the New England District ACOE, which is provided in Appendix F.

For basic information on the use of soils in wetland delineation see “The Buzzards Bay National Estuary Program Pocket Guide to Delineating Wetlands.”

¹ We have included information from the 2013 Errata for this document.

DEP Short List

The DEP list of seven most common wetland soils is found on Page 29 of the DEP Manual.

These seven hydric soil morphologies are not an exhaustive or exclusive list. DEP refers to them as “some hydric soil indicators.”

Please note that the federal list in the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0) has a slightly different take on Histosols and the criteria for a depleted matrix.

These differences are described in the “DEP Particularly Difficult Soils” section.

DEP Short List: 1. Histosols (organic soils)

Histosols are soils with at least 16 inches of organic material measured from the soil surface.



Photo Credit: Jim Turenne

BBNEP note: This soil is not typically found at the BVW edge.

DEP Short List: 2. Histic Epipedons

These are soils with 8 to 16 inches of organic material measured from the soil surface.



Photo Credit: Field Indicators of Hydric Soils in the United States v. 6

DEP Short List: 3. Sulfidic Material.

A strong "rotten egg" smell generally is noticed immediately after the soil test hole is dug.



Photo credit: Slide from NRCS, Wetland Science Institute, Power Point Presentation "Field Indicators of Hydric Soils" available at <http://www.maenvirothon.org/hydric%20soils.ppt>

BBNEP note: This soil is associated with salt marshes and is not typically found at the BVW edge.

DEP Short List: 4. Gleyed Matrix

Soils that are predominantly neutral gray, or occasionally greenish or bluish gray in color within 12 inches from the bottom of the 0-horizon. (The Munsell Soil Color Charts have special pages for gleyed soils.)



Photo Credit: Field Indicators of Hydric Soils in the United States v. 6 This soil has a gleyed matrix in the lowest layer, starting about 7 in. from the soil surface. The layer above the gleyed matrix has a depleted matrix.

DEP Short List: 4. Gleyed soils, cont.



For hydric soil determinations, a gleyed matrix has the hues and chroma identified in this illustration with a value of 4 or more. [Due to inaccurate color reproduction, do not use this page to determine soil colors in the field.]

DEP Short List: 5. Depleted Matrix (1)

Soils with a matrix chroma of 0 or 1 and values of 4 or higher within 12 inches from the bottom of the 0-horizon



0a, 7.5YR 2/0,
muck.

A, 10YR 2/1,
mucky silt loam

2Cg, 2.5Y 6/1,
silt loam

Typical profile of a Birdsall mucky silt loam soil. Birdsall soils are very poorly drained soils formed in water-laid deposits of silt and very fine sand.

Photo Credit: Peter Fletcher

DEP Short List: 6. Depleted Matrix (2)

Within 12 inches from the bottom of the 0-horizon, soils with a chroma of 2 or less and values of 4 or higher in the matrix, and mottles with a chroma of 3 or higher.



Photo from: NRCS, Wetland Science Institute, Power Point Presentation
“Field Indicators of Hydric Soils” available at
<http://www.maenvirothon.org/hydric%20soils.ppt>

DEP Short List: 7. Redox Depletions

Within 12 inches from the bottom of the 0-horizon, soils with a matrix chroma of 3 and values of 4 or higher, with 10 percent or more low-chroma mottles, as well as indicators of saturation (i.e., mottles, oxidized rhizospheres, concretions, nodules) within 6 inches of the soil surface.



Photo from: “Redoximorphic Features” presentation developed by Michael Whited, NRCS - Wetland Science Inst. August, 2000

<http://www.maenvirothon.org/hydric%20soils.ppt>

BBNEP Note: DEP Particularly Difficult Soil Indicators XI. A. Redox Depletions and XI. B. Redox Depletions & Thick Very Dark Surface.

DEP Difficult Soils

On pages 30 and 31 of the DEP Delineation Manual, DEP lists six soil types that are “difficult to analyze.”

Most of these soils have suggested hydric morphologies that are listed in the following pages. Where additional soil morphologies for each type are listed in the [Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region \(Version 2.0\)](#), [Field Indicators of Hydric Soils in the United States, Version 7](#) (including the 2013 Errata), or [Field Indicators for Identifying Hydric Soils in New England, Version 3](#), a reference to them is provided.



Photo credit: John Rockwell, Soil samples from Carver. Wettest from right to left. These are disturbed, sandy spodosols (evergreen forest soils). Notice the redoximorphic features in all samples.

DEP Difficult Soils: Sandy Soils

“Sandy soils” refers to soil materials with a USDA soil texture of loamy fine sand and coarser.

Soil colors often are not distinctive in most sandy soils. Instead, look for these indicators of hydric sandy soils:

- a) high organic content in the surface layer (typically darker colors with values less than 3 and chroma of 2 or less) with mottles or other indicators of saturation directly below;
- b) organic streaking directly below the A-horizon; or
- c) matrix chroma of 3 (from the Munsell Soil Color Charts) in the top 12 inches of soil measured from the bottom of the O-horizon, with distinct or prominent mottling.



Photo credit: Southeast Soil & Water Service, <http://www.hydricsoils.com>

Note: Indicators of hydric soils may be lacking altogether in the soil of newly formed sand bars and interdunal depressions.

BBNEP Note: See the “S” Indicators and X. Sandy with Redox and Thick, Very Dark Surface, in the “DEP Particularly Difficult Soils” section for more sandy soils hydric morphologies.

DEP Difficult Soils: Floodplain Soils

These soils usually are characterized by distinctly layered soil material. The layers form when new sediment is deposited during flood events. As a result of this pattern of deposition, hydric soil indicators may never form, or may be buried even though saturated or inundated conditions are present long enough to create wetland hydrology.




BBNEP Note: DEP has not provided any morphological criteria for this difficult soil. See “DEP Particularly Difficult Soils” Indicators A5, F12, and F19.

DEP Difficult Soils: Soil from Highly Colored Parent Material

Some soils derived from highly colored parent material have strong red, brown, or black colors. As a result, the gray colors indicative of hydric soils may not be obvious. Red soils generally are confined to certain areas within the Connecticut River Valley. Brown soils derived from Brimfield schists generally are found in and around the town of Brimfield. Black soils generally are confined to southeastern Massachusetts (principally Bristol County).

Dark Parent Material




A-horizons greater than or equal to 12 inches thick with values less than 3 and chroma of 2 or less are difficult to analyze because indicators of saturation are difficult to see. Therefore, look directly below the A-horizon for a matrix chroma of 1 or less and values of 4 or higher.

If the matrix color directly below the thick and dark A-horizon is chroma 2 and value 4 or higher, other indicators of saturation need to be present in the soil directly below the A-horizon. In uncommon situations, it may be necessary to dig deeper to evaluate colors below the A-horizon.

BBPNEP Note: Also see “DEP Particularly Difficult Soils” Indicators A11, A12, and F6. In addition to the Regional Supplement soils, see Indicator VIII Dark Soil Materials

DEP Difficult Soils: Soil from Highly Colored Parent Material, cont.

Red Parent Material



These maps were prepared by Al Averill and Darlene Monds of the NRCS Massachusetts office.

BBNEP Note: DEP has not provided any morphological criteria for this difficult soil.. For Red Parent Materials see “DEP Particularly Difficult Soils” Indicator F21 and TF2. These difficult soils are not found in the Buzzards Bay watershed

DEP Difficult Soils: A-horizons that are Thick and Very Dark.

A-horizons greater than or equal to 12 inches thick with values less than 3 and chroma of 2 or less are difficult to analyze because indicators of saturation are difficult to see. Therefore, look directly below the A-horizon for a matrix chroma of 1 or less and values of 4 or higher. If the matrix color directly below the thick and dark A-horizon is chroma 2 and value 4 or higher, other indicators of saturation need to be present in the soil directly below the A-horizon. In uncommon situations, it may be necessary to dig deeper to evaluate colors below the A-horizon.



Photo credit: NRCS

BBPNEP Note: Also see “DEP Particularly Difficult Soils” Indicators A11, A12, F6, X.C. Sandy with Redox & Thick, Very Dark Surface, XI. B. Redox Depletions & Thick Very Dark Surface, and VII, Dark Surface Materials.

DEP Difficult Soils Evergreen Forest Soils

Sandy soils on Cape Cod and other areas dominated by evergreen trees may possess gray colored E-horizons just beneath the surface. These colors are not necessarily the result of saturation or inundation, but form as a result of the leaching of organic material and aluminum and iron oxides by organic acids. These soils are called **spodosols** and the gray layer that forms below the surface is known as the E-horizon. Organic material and aluminum and iron oxides are deposited in a layer below the E-horizon called the spodic horizon.

Hydric indicators in spodosols include a combination of two or more of the following features, with one occurring within the upper 12 inches of the soil surface and others documented below the soil surface:

- a) a thick, black, sandy surface layer;
- b) organic streaking in the E-horizon;
- c) mottles within the E-horizon;
- d) oxidized rhizospheres within the A or E-horizon;
- e) iron concretions/nodules within the E-horizon or spodic horizon;
- f) a partially or wholly cemented spodic horizon usually within 18 inches of the surface measured from the bottom of the O-horizon; and mottling within the spodic horizon.

Non-hydric spodosols can be recognized by brightly colored soil material below the E-horizon and without mottles or other indicators of saturation.



Photo Credit: John Rockwell. Attendees of the MACC/BBNEP 2009 Advanced session check colors for a partially cemented spodic horizon.

DEP Difficult Soils Evergreen Forest Soils, cont.

BBPNP Note: These soils can be found throughout the Buzzards Bay watershed. They can be found in loams as well as the more often occurring sandy soils. See Appendix C for information on spodosol field identification.

See also “DEP Particularly Difficult Soils” Indicators S8, S9, TA6, IX. A., B., & C. Spodosols.



Photo Credit: John Rockwell. These soils are quite varied but were all found within 100 feet of each other at Washburn Park in Marion, Massachusetts. The three on the right are samples from a disturbed site. The left sample is a wet spodosol.

DEP Difficult Soils: Areas where the Hydrology has been Recently Altered.

In areas where the hydrology has been recently altered, hydric soil indicators may not accurately reflect the current hydrology of the site. Areas that have been recently flooded - or where the water table has risen due to flooding or some other change in hydrologic conditions - may not exhibit hydric soil characteristics. These areas may not have been saturated long enough to develop hydric characteristics. Conversely, areas that have been effectively drained and wetland hydrology is no longer present may still possess hydric soil indicators. Where there is evidence that the hydrology has been substantially altered at a site, careful evaluation of vegetation, soils, and other indicators of hydrology should be made before making a final delineation. Altered areas are particularly difficult to evaluate and require special attention.

Photo credit: Muncie Sanitary District



BBNEP Note: Use Google Earth to determine the timing of the alteration. The area may be subject to enforcement action depending on the date of the alteration activity.

DEP Particularly Difficult Soils

In the first paragraph on page 30 of the DEO Delineation Manual, DEP states, “In particularly difficult cases, consultation with the Natural Resources Conservation Service (NRCS) is recommended.”

NRCS uses the latest federal hydric soil list. A list of these soils can be found in the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0), which is based on Field Indicators of Hydric Soils in the United States, Version 7 (updated by 2013 Erratta).

Many of these hydric soil indicators were developed specifically for wetland delineation purposes. During the development of these indicators, soils in the interior of wetlands were not always examined; therefore, there are wetlands that lack any of the approved hydric soil indicators in the wettest interior portions.

Wetland delineators and other users of these hydric soil indicators should concentrate their sampling efforts near the wetland edge and, if these soils are hydric, assume that soils in the wetter, interior portions of the wetland are also hydric, even if they lack an indicator.

Hydric soil indicators are presented in three groups. Indicators for “All Soils” are used in any soil regardless of texture. Indicators for “Sandy Soils” are used in soil layers with USDA textures of loamy fine sand or coarser. Indicators for “Loamy and Clayey Soils” are used with soil layers of loamy very fine sand and finer.

Both sandy and loamy/clayey layers may be present in the same soil profile. Therefore, a soil that contains a loamy surface layer over sand is hydric if it meets all of the requirements of matrix color, amount and contrast of redox concentrations, depth, and thickness for a specific A (All Soils), F (Loamy and Clayey Soils), or S (Sandy Soils) indicator.

It is permissible to combine certain hydric soil indicators if all requirements of the individual indicators are met except thickness (see Hydric Soil Technical Note 4, http://soils.usda.gov/use/hydric/ntchs/tech_notes/index.html). See Appendix B for notes on thickness.

“All soils” refers to soils with any USDA soil texture. Use the “A” indicators regardless of soil texture.

“Sandy soils” refers to soil materials with a USDA soil texture of loamy fine sand and coarser. Use the “S” indicators in soil layers consisting of sandy soil materials.

“Loamy and clayey soils” refers to soil materials with USDA textures of loamy very fine sand and finer. Use the “F” indicators in soil layers consisting of loamy or clayey soil materials

All mineral layers above any of the layers meeting an A, S, or F indicator must have a dominant chroma of 2 or less, or the layer(s) with a dominant chroma of more than 2 must be less than 6 in. thick, except for indicators S6, F8, F12, and F21 to meet any hydric soil indicator. Nodules and concretions are not considered to be redox concentrations unless otherwise noted.

In problem areas it is permissible to use soil morphologies described in Field Indicators for Hydric Soils in New England, Version 3, (See memo from Ruth Ladd in Appendix F). This is particularly useful in areas with red soil materials and in areas with spodosols.

Some of these soil morphologies should not be considered conclusive evidence of saturated or inundated conditions without a independent confirmation of hydrology. These soils are so noted in the BBNEP Notes.

A list of wetland hydrology indicators can be found in the Postscript on page 65.

DEP Particularly Difficult Soils: Indicator A1: Histosol

In most Histosols, 16 in. or more of the upper 32 in. is organic soil material. Histosols also include soils that have organic soil material of any thickness over rock or fragmental soil material that has interstices filled with organic soil material. Organic soil material has an organic carbon content (by weight) of 12 to 18 percent or more, depending on the clay content of the soil. The material includes muck (sapric soil material), mucky peat (hemic soil material), or peat (fibric soil material).



This Histosol consists of only a few inches of organic soil material over bedrock in a shallow glacial groove.

BBNEP Note: Indicator A1 differs from DEP Short List 1. Histosols. To meet the Indicator A1 criteria, Histosols may be thinner over bedrock areas than the 16 inches called for in the DEP Short List definition.

DEP Particularly Difficult Soils: Indicator A5: Stratified Layers

Several stratified layers starting within 6 in. of the soil surface. At least one of the layers has a value of 3 or less with a chroma of 1 or less or it is muck, mucky peat, peat, or mucky modified mineral texture. The remaining layers have chromas of 2 or less (see figure below).

Any sandy material that constitutes the layer with a value of 3 or less and a chroma of 1 or less, when viewed with a 10- or 15-power hand lens, must have at least 70 percent of the visible soil particles masked with organic material (see figure on next page). When viewed without a hand lens, the material appears to be nearly 100 percent masked.



Stratified layers in loamy material.

DEP Particularly Difficult Soils: Indicator A5: Stratified Layers, cont.



Stratified layers in sandy material

User Notes: Use of this indicator may require assistance from a soil scientist with local experience. An undisturbed sample must be observed. Individual strata are dominantly less than 1 in. thick.

A hand lens can aid in the identification of this indicator.

Many alluvial soils have stratified layers at depths greater than 6 in.; these do not fit this indicator. Many alluvial soils have stratified layers at the required depths but lack a chroma of 2 or less; these do not fit this indicator.

Stratified layers occur in any type of soil material, generally in floodplains and other areas where wet soils are subject to rapid and repeated burial with thin deposits of sediment.

DEP Particularly Difficult Soils: Indicator A11: Depleted Below Dark Surface

A layer with a depleted or gleyed matrix that has 60 percent or more chroma of 2 or less, starting within 12 in. of the soil surface, and having a minimum thickness of either:

- 6 in. or
- 2 in. if the 2 in. consists of fragmental soil material.

Loamy/clayey layer(s) above the depleted or gleyed matrix must have a value of 3 or less and chroma of 2 or less. Any sandy material above the depleted or gleyed matrix must have a value of 3 or less and chroma of 1 or less and, when viewed with a 10- or 15-power hand lens, must have at least 70 percent of the visible soil particles masked with organic material. When viewed without a hand lens, the material appears to be nearly 100 percent masked.

User Notes: This indicator often occurs in hydric soils that have dark-colored surface layers, such as umbric epipedons and dark-colored ochric epipedons (see figure below).

For soils that have dark surface layers greater than 12 in. thick, use indicator A12.




In this soil, a depleted matrix starts immediately below the black surface layer at approximately 11 in. (28 cm).

DEP Particularly Difficult Soils: Indicator A11: Depleted Below Dark Surface, cont.

Two percent or more distinct or prominent redox concentrations, including iron/manganese soft masses, pore linings, or both, are required in soils that have matrix values/chromas of 4/1, 4/2, and 5/2 (see figure below). If the soil is saturated at the time of sampling, it may be necessary to let it dry to a moist condition for redox features to become visible.

Illustration of values and chromas that require 2 percent or more



distinct or prominent redox concentrations and those that do not, for hue 10YR, to meet the definition of a depleted matrix. Due to inaccurate color reproduction, do not use this page to determine soil colors in the field.

BBNEP Note: This depleted criteria differs from DEP Short List #6 Depleted Matrix as values/chromas of 6/2, 7/2, and 8/2 do not require redox features.

DEP Particularly Difficult Soils: Indicator A12: Thick Dark Surface

Technical Description: A layer at least 6 in. thick with a depleted or gleyed matrix that has 60 percent or more chroma of 2 or less starting below 12 in. of the surface. The layer(s) above the depleted or gleyed matrix must have a value of 2.5 or less and chroma of 1 or less to a depth of at least 12 in. and a value of 3 or less and chroma of 1 or less in any remaining layers above the depleted or gleyed matrix. Any sandy material above the depleted or gleyed matrix, when viewed with a 10- or 15-power hand lens, must have at least 70 percent of the visible soil particles masked with organic material. When viewed without a hand lens, the material appears to be nearly 100 percent masked.

User Notes: The soil has a depleted matrix or gleyed matrix below a black or very dark gray surface layer 12 in. or more thick (see figure below). This indicator is most often associated with overthickened soils in concave landscape positions. Two percent or more distinct or prominent redox concentrations, including iron/manganese soft masses, pore linings, or both, are required in soils that have matrix values/chromas of 4/1, 4/2, and 5/2 (see figure on p 25). If the soil is saturated at the time of sampling, it may be necessary to let it dry to a moist condition for redox features to become visible.



Deep observations may be necessary to identify the depleted or gleyed matrix below a thick, dark surface layer. In this example, the depleted matrix starts at 20 in.

BBNEP Note: For thick surface horizons see also Indicators F6, X.C. Sandy with Redox & Thick, Very Dark Surface, XI. B. Redox Depletions & Thick Very Dark Surface, and VII, Dark Surface Materials.

DEP Particularly Difficult Soils: Indicator S1: Sandy Mucky Mineral

Technical Description: A layer of mucky modified sandy soil material 2 in. or more thick starting within 6 in. of the soil surface (see figure below).

User Notes: This indicator is uncommon but is found in localized areas in this region. *Mucky* is a USDA texture modifier for mineral soils. The organic carbon content is at least 5 percent and ranges up to 14 percent for sandy soils. The percentage requirement is dependent upon the clay content of the soil; the higher the clay content, the higher the organic carbon requirement. See the glossary of *Field Indicators of Hydric Soils in the United States* (USDA Natural Resources Conservation Service 2010) for the definition of mucky modified mineral texture. A field procedure for identifying mucky mineral soil material is presented in Appendix D.



The mucky modified sandy layer is approximately 3 in. thick. Scale in inches on the right side of ruler.

DEP Particularly Difficult Soils: Indicator S6: Stripped Matrix

Technical Description: A layer starting within 6 in. of the soil surface in which iron/manganese oxides and/or organic matter have been stripped from the matrix and the primary base color of the soil material has been exposed. The stripped areas and translocated oxides and/or organic matter form a faintly contrasting pattern of two or more colors with diffuse boundaries. The stripped zones are 10 percent or more of the volume and are rounded.

User Notes: This indicator includes the indicator previously named streaking (Environmental Laboratory 1987). The stripped areas are typically 0.5 to 1 in. in size but may be larger or smaller. Commonly, the stripped areas have a value of 5 or more and chroma of 1 and/or 2 and unstripped areas have a chroma of 3 and/or 4 (see figure below).

However, there are no specific color requirements for this indicator. The mobilization and translocation of the oxides and/or organic matter are the important processes involved in this indicator and should result in splotchy coated and uncoated soil areas. A 10-power hand lens can be helpful in seeing stripped and unstripped areas. This may be a difficult pattern to recognize and is often more evident in a horizontal slice.

This is a very common indicator of hydric soils and is often used to identify the hydric/non-hydric boundary in sandy soils. This indicator is found in all wetland types and all wet landscape positions.



In this example, a faint splotchy pattern of stripped and unstripped areas lies beneath a thin dark surface layer.

DEP Particularly Difficult Soils: Indicator S7: Dark Surface

Technical Description: A layer 4 in. thick starting within 6 in. of the soil surface with a matrix value of 3 or less and chroma of 1 or less. When viewed with a 10- or 15-power hand lens, at least 70 percent of the visible soil particles must be masked with organic material. When viewed without a hand lens, the material appears to be nearly 100 percent masked. The matrix color of the layer immediately below the dark layer must have the same colors as those described above or any color that has a chroma of 2 or less.

User Notes: If the dark layer is greater than 4 in. thick, then the indicator is met, because any dark soil material in excess of 4 in. meets the requirement that “the layer immediately below the dark layer must have the same colors as those described above” If the dark layer is exactly 4 in. thick, then the material immediately below must have a matrix chroma of 2 or less.

This indicator is applicable to interdunal swales along the Atlantic Ocean. The organic carbon content of this indicator is slightly less than that required for “mucky.” An undisturbed sample must be observed (see figure below). Many moderately wet soils have a ratio of about 50 percent of soil particles covered or coated with organic matter to about 50 percent uncoated or uncovered soil particles, giving the soil a salt-and-pepper appearance. Where the percent coverage by organic matter is less than 70 percent, the Dark Surface indicator is not present.



Example of Indicator S7 (Dark Surface) in a sandy soil. Scale in inches on right.

DEP Particularly Difficult Soils: Indicator S8: Polyvalue Below Surface

Technical Description: A layer with a value of 3 or less and chroma of 1 or less starting within 6 in. of the soil surface. When viewed with a 10- or 15-power hand lens, at least 70 percent of the visible soil particles in this layer must be masked with organic material. When viewed without a hand lens, the material appears to be nearly 100 percent masked. Immediately below this layer, 5 percent or more of the soil volume has a value of 3 or less and chroma of 1 or less and the remainder of the soil volume has a value of 4 or more and chroma of 1 or less to a depth of 12 in. or to the spodic horizon, whichever is less.

User Notes: This indicator applies to soils with a very dark gray or black surface or near-surface layer that is underlain by a layer in which organic matter has been differentially distributed within the soil by water movement (see figure below). The mobilization and translocation of organic matter result in splotchy coated and uncoated soil areas, as described in the Sandy Redox (S5) and Stripped Matrix (S6) indicators, except that for S8 the whole soil is in shades of black and gray. The chroma of 1 or less is critical because it limits application of this indicator to only those soils that are depleted of iron. This indicator includes the indicator previously termed “streaking.” See Appendix C for field criteria to identify a spodic horizon.



In this soil, the splotchy pattern below the dark surface is due to mobilization and translocation of organic matter. Scale in inches.

BBNEP Note: For additional hydric spodosol morphologies see Indicators, S9, TA6, IX. A., B., & C. Spodosols.

DEP Particularly Difficult Soils: Indicator S9: Thin Dark Surface

Technical Description: A layer 2 in. or more thick starting within the upper 6 in of the soil, with a value of 3 or less and chroma of 1 or less. When viewed with a 10- or 15-power hand lens, at least 70 percent of the visible soil particles in this layer must be masked with organic material. When viewed without a hand lens, the material appears to be nearly 100 percent masked. This layer is underlain by a layer(s) with a value of 4 or less and chroma of 1 or less to a depth of 12 in. (30 cm) or to the spodic horizon, whichever is less.

User Notes: This indicator applies to soils with a very dark gray or black near-surface layer that is at least 2 in. thick and is underlain by a layer in which organic matter has been carried downward by flowing water (see figure below). The mobilization and translocation of organic matter result in an even distribution of organic matter in the eluvial (E) horizon. The chroma of 1 or less is critical because it limits application of this indicator to only those soils that are depleted of iron. This indicator commonly occurs in hydric Spodosols; however, a spodic horizon is not required. See Appendix C for field criteria to identify a spodic horizon.



Example of Indicator S9 (Thin Dark Surface). Scale in inches on right.

BBNEP Note:
For additional hydric spodosol morphologies see Indicators S8, TA6, IX A., B., & C. Spodosols.

DEP Particularly Difficult Soils: Indicator F3: Depleted Matrix

Technical Description: A layer that has a depleted matrix with 60 percent or more chroma of 2 or less and that has a minimum thickness of either:

- 2 in. if the 2 in. is entirely within the upper 6 in. of the soil, or
- 6 in. starting within 10 in. of the soil surface.



Example of indicator F3 (Depleted Matrix), in which redox concentrations extend nearly to the surface.

BBNEP Note: This depleted criteria differs from DEP Short List #6 Depleted Matrix as values/chromas of 6/2, 7/2, and 8/2 do not require redox features.



Illustration of values and chromas that require 2 percent or more distinct or prominent redox concentrations and those that do not, for hue 10YR, to meet the definition of a depleted matrix. Due to inaccurate color reproduction, do not use this page to determine soil colors in the field. Background image from the Munsell Soil Color Charts reprinted courtesy of Munsell Color Services Lab, a part of X-Rite, Inc. (Gretag/Macbeth 2000).

DEP Particularly Difficult Soils: Indicator F6: Redox Dark Surface

Technical Description: A layer that is at least 4 in. thick, is entirely within the upper 12 in. of the mineral soil, and has a:

- matrix value of 3 or less and chroma of 1 or less and 2 percent or more distinct or prominent redox concentrations occurring as soft masses or pore linings, or
- matrix value of 3 or less and chroma of 2 or less and 5 percent or more distinct or prominent redox concentrations occurring as soft masses or pore linings.

User Notes: This is a very common indicator used to delineate wetlands. Redox concentrations are often small and difficult to see in mineral soils that have dark (value of 3 or less) surface layers due to high Organic-matter content (see figure below). The organic matter masks some or all of the concentrations that may be present; it also masks the diffuse boundaries of the concentrations and makes them appear to be more sharp. Careful examination is required to see what are often brownish redox concentrations in the darkened materials. If the soil is saturated at the time of sampling, it may be necessary to let it dry at least to a moist condition for redox features to become visible. In some cases, further drying of the samples makes the concentrations (if present) easier to see. A hand lens may be helpful in seeing and describing small redox concentrations. Care should be taken to examine the interior of soil peds for redox concentrations. Dry colors, if used, also must have matrix chromas of 1 or 2, and the redox concentrations must be distinct or prominent. For soils with thick, dark surface layers, see also indicators A11 (Depleted Below Dark Surface) and A12 (Thick Dark Surface).

In soils that are wet because of subsurface saturation, the layer immediately below the dark epipedon will likely have a depleted or gleyed matrix (see the Glossary for definitions). Soils that are wet because of ponding or have a shallow, perched layer of saturation may not always have a depleted/gleyed matrix below the dark surface. This morphology has been observed in soils that have been compacted by tillage and other means. It is recommended that delineators evaluate the hydrologic source and examine and describe the layer below the dark-colored epipedon when applying this indicator.



BBPNP Note: Also see “DEP Particularly Difficult Soils” Indicators A11, A12, X.C. Sandy with Redox & Thick, Very Dark Surface, XI. B. Redox Depletions & Thick Very Dark Surface, and VII, Dark Surface Materials.

R

edox features can be small and difficult to see within a dark soil layer.

DEP Particularly Difficult Soils: Indicator F7: Depleted Dark Surface

Technical Description: Redox depletions with a value of 5 or more and chroma of 2 or less in a layer that is at least 4 in. thick, is entirely within the upper 12 in. of the mineral soil (see figure below), and has a:

- matrix value of 3 or less and chroma of 1 or less and 10 percent or more redox depletions, or
- matrix value of 3 or less and chroma of 2 or less and 20 percent or more redox depletions.

User Notes: Care should be taken not to mistake the mixing of eluvial (leached) layers that have high value and low chroma (E horizon) or Illuvial layers that have accumulated carbonates (calcic horizon) into the surface layer as depletions. Mixing of layers can be caused by burrowing animals or cultivation. Pieces of deeper layers that become incorporated into the surface layer are not redox depletions. Knowledge of local conditions is required in areas where light-colored eluvial layers and/or layers high in carbonates may be present. In soils that are wet because of subsurface saturation, the layer immediately below the dark surface is likely to have a depleted or gleyed matrix. Redox depletions are usually associated with microsites that have redox concentrations occurring as pore linings or masses within the depletion(s) or surrounding the depletion(s).




Redox depletions (lighter colored areas) are scattered within the darker matrix. Scale is in centimeters.

DEP Particularly Difficult Soils: Indicator X.C. Sandy With Redox & Thick, Very Dark Surface

Technical Description: Soils that do not have a *spodic* horizon and beginning within 15 inches of the *top of the mineral soil material* and directly underlying a *thick, very dark A* or *Ap horizon* there is a horizon with a *loamy fine sand or coarser texture* with a *matrix color* due to wetness of chroma 3 or less, value 4 or more, with 2 percent or more *redoximorphic features*.

User Notes: Field investigations have documented some situations where the conditions for a hydric soil are present and the matrix chroma directly underlying the *A* or *Ap horizon* is greater than 3. These soil conditions are associated with iron-enriched groundwater discharge areas. These soils are considered problem soils. When soil morphology seems inconsistent with the landscape, vegetation, or observable hydrology, the assistance of an experienced soil or wetland scientist may be needed to determine whether the soil is hydric.

Terms or phrases in italics are defined in the Glossary of Terms of Field Indicators for Identifying Hydric Soils in New England, Version 3.




BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator XI. B. Redox Depletions & Thick Very Dark Surface (Any Texture)

Technical Description: Soils that do not have a *spodic horizon* and beginning within 15 inches of the *top of the mineral soil material* and directly underlying a *thick, very dark Ap horizon*, is a horizon with 5 percent or more *redox depletions* and within 20 inches of the *top of the mineral soil material* there is a horizon with a *depleted or gleyed matrix* (for soils with *moderate to strong structure*, the *matrix color* and percent *redox depletions* are recorded for *ped interiors*).

Terms or phrases in italics are defined in the Glossary of Terms of Field Indicators for Identifying Hydric Soils in New England, Version 3.



BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator F8: Redox Depressions

Technical Description: In closed depressions subject to ponding, 5 percent or more distinct or prominent redox concentrations occurring as soft masses or pore linings in a layer that is 2 in. or more thick and is entirely within the upper 6 in. of the soil (see figure below).

User Notes: This indicator occurs on depressional landforms, such as vernal pools and potholes, but not microdepressions on convex landscapes. Closed depressions often occur within flats or floodplain landscapes. *Note that there is no color requirement for the soil matrix.* The layer containing redox concentrations may extend below 6 in. as long as at least 2 in. occurs within 6 in. of the surface. If the soil is saturated at the time of sampling, it may be necessary to let it dry to a moist condition for redox features to become visible.

This is a common but often overlooked indicator found at the wetland/non-wetland boundary on depressional sites.




In this example, the layer of redox concentrations begins at the soil surface and is slightly more than 2 in. thick.

DEP Particularly Difficult Soils: Indicator A10: 2 cm Muck

Technical Description: A layer of muck 0.75 in. (2 cm) or more thick with a value of 3 or less and chroma of 1 or less, starting within 6 in. of the soil surface.

Applicable Subregions: Long Island/Cape Cod (MLRA 149B of LRR S).

User Notes: Normally the muck layer is at the soil surface; however, it may occur at any depth within 6 in. (15 cm) of the surface. Muck is Sapric soil material with at least 12 to 18 percent organic carbon. Organic soil material is called muck if virtually all of the material has undergone sufficient decomposition to limit recognition of the plant parts. Hemic (mucky peat) and fibric (peat) soil materials do not qualify. To determine if muck is present, first remove loose leaves, needles, bark, and other easily identified plant remains. This is sometimes called leaf litter, a duff layer, or a leaf or root mat. Then examine for decomposed organic soil material. Generally, muck is black and has a greasy feel; sand grains should not be. Determination of this indicator is made below the leaf or root mat; however, root mats that meet the definition of hemic or fibric soil material are included in the decisionmaking process for indicators A1 (Histosol) and A2 (Histic Epipedon).



This soil has more than 2 cm of muck, starting at 8 cm on the left measuring tape. Photo credit: Field Indicators of Hydric Soils in the United States, A Guide for Identifying and Delineating Hydric Soils, Version 7.0, 2010

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator A16: Coast Prairie Redox

Technical Description: A layer starting within 6 in. of the soil surface that is at least 4 in. (10 cm) thick and has a matrix chroma of 3 or less with 2 percent or more distinct or prominent redox concentrations occurring as soft masses and/or pore linings.

Applicable Subregions: **not** in the Long Island/Cape Cod Subregion (MLRA 149B of LRR S).

User Notes: These hydric soils occur mainly on depressional and intermound landforms. Redox concentrations occur mainly as iron- dominated pore linings. Common to many redox concentrations are required. If the soil is saturated at the time of sampling, it may be necessary to let it dry to a moist condition for redox features to become visible. Chroma 3 matrices are allowed because they may be the color of stripped sand grains, or because few to common sand-sized reddish particles may be present and may prevent obtaining a chroma of 2 or less.

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator S3: 5 cm Mucky Peat or Peat

Technical Description: A layer of mucky peat or peat 2 in. or more thick with a value of 3 or less and chroma of 2 or less, starting within 6 in. of the soil surface, and underlain by sandy soil material.

Applicable Subregions: not in the Long Island/Cape Cod Subregion (MLRA 149B of LRR S).

User Notes: In this region, this indicator is applicable primarily to interdunal swales along the Great Lakes and Atlantic coast. Mucky peat (hemic soil material) and peat (fibric soil material) have at least 12 to 18 percent organic carbon. Organic soil material is called peat if virtually all of the plant remains are sufficiently intact to permit identification of plant remains. Mucky peat is an intermediate stage of decomposition between peat and highly decomposed muck. See the glossary of Field Indicators of Hydric Soils in the United States (USDA Natural Resources Conservation Service 2010) for definitions. See the Concepts section of this chapter for field methods to identify organic soil materials.

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator F12: Iron-Manganese Masses

Technical Description: On floodplains, a layer 4 in. or more thick with 40 percent or more chroma of 2 or less and 2 percent or more distinct or prominent redox concentrations occurring as soft iron-manganese masses with diffuse boundaries. The layer occurs entirely within 12 in. of the soil surface. Iron-manganese masses have a value and chroma of 3 or less. Most commonly, they are black. The thickness requirement is waived if the layer is the mineral surface layer.

Applicable Subregions: not in the Long Island/Cape Cod Subregion (MLRA 149B of LRR S).

User Notes: These iron-manganese masses generally are small (2 to 5 mm in size) and have value and chroma of 3 or less. They can be dominated by manganese and, therefore, have a color approaching black (see figure below). If the soil is saturated at the time of sampling, it may be necessary to let it dry to a moist condition for redox features to become visible. The low matrix chroma must be the result of wetness and not be a relict or parent material feature. Iron-manganese masses should not be confused with the larger and redder iron nodules associated with plinthite or with concretions that have sharp boundaries. This indicator occurs on floodplains such as those of the Mississippi, Hudson, and Penobscot Rivers.



Iron-manganese masses (black spots) in a 40 percent depleted matrix. Scale is in inches.

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator F19: Piedmont Floodplain Soils

Technical Description: On floodplains, a mineral layer at least 6 in. thick starting within 10 in. of the soil surface with a matrix (60 percent or more of the volume) chroma of less than 4 and 20 percent or more distinct or prominent redox concentrations occurring as soft masses or pore linings.

Applicable Subregions: For use with problem soils in the Long Island/Cape Cod Subregion (MLRA 149B of LRR S) (See Appendix A).

User Notes: This indicator is for use or testing on flood plains in the Mid-Atlantic and Southern Piedmont Provinces and areas where sediments derived from the Piedmont have been deposited on flood plains on the Coastal Plain. This indicator does not apply to stream terraces, which are associated with a previous stream level and are representative of an abandoned flood plain. While these soils are found on flood plains, flooding may be rare and groundwater is often the source of hydrology.



The Piedmont Floodplain Soils indicator is restricted to floodplains that are actively receiving sediments and groundwater discharge with high iron content. Photo by M. Rabenhorst. Scale in 4-in. increments.


BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Addition indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator F21 Red Parent Materials

Technical Description: A layer derived from red parent materials that is at least 4 inches thick, starting within 10 inches of the soil surface with a hue of 7.5YR or redder. The matrix has a value and chroma greater than 2 and less than or equal to 4. The layer must contain 10 percent or more depletions and/or distinct or prominent redox concentrations occurring as soft masses or pore linings. Redox depletions should differ in color by having:

- Value one or more higher and chroma one or more lower than the matrix, or
- Value of 4 or more and chroma of 2 or less.

User Notes: This indicator was developed for use in areas of red parent material, such as residuum in the Piedmont Province Triassic lowlands section or the Paleozoic “red beds” of the Appalachian Mountains, and in alluvium or colluvium derived from these materials. In glaciated areas, the indicator may form in glacial till, outwash, deltaic sediments, or glaciolacustrine sediments derived from similar red lithologies. In order to confirm that it is appropriate to apply this indicator to particular soils, soils formed from similar parent materials in the area should have been evaluated to determine their Color Change Propensity Index (CCPI) and be shown to have CCPI values below 30 (Rabenhorst and Parikh, 2000.) It cannot be assumed that sediment overlying red colored bedrock is derived solely from that bedrock. The total percentage of all redox concentrations and redox depletions must add up to at least 10% to meet the threshold for this indicator.



This indicator should be used only in areas of red parent material that is resistant to reduction. Not all red soils formed in red parent material.

DEP Particularly Difficult Soils: Indicator F21 Red Parent Materials, cont.

This indicator is typically found at the boundary between hydric and non-hydric soils. Users that encounter a depleted matrix in the upper part should consider F3-Depleted Matrix. F3 is often found in sites that are anaerobic for a longer period. Users that encounter a dark soil surface (value 3 or less and chroma 2 or less) should consider F6-Redox Dark Surface or F7-Depleted Dark Surface. If the site is in a closed depression subject to ponding users should consider F8-Redox Depressions.



Indicator F3 (Depleted Matrix) in red parent material. If a soil that formed in red parent material stays wet and anaerobic long enough, it may develop the characteristics of DEP Particularly Difficult Soils: Indicator F3: Depleted Matrix.

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement. Also see Indicator TF2 Red Parent Material (next page). Questions about hydric soil morphologies in red parent materials should be directed to Donald Parizek, Soil Scientist at the Connecticut NRCS office in Tolland.

DEP Particularly Difficult Soils: Indicator TF2 Red Parent Materials

Technical Description: In parent material with hue of 7.5YR or redder, a layer at least 4 inches thick with a matrix value and chroma of 4 or less and 2 percent or more redox depletions and/or redox concentrations occurring as soft masses and/or pore linings. The layer is entirely within 12 inches of the soil surface. The minimum thickness requirement is 2 inches if the layer is the mineral surface layer.

User Notes: This indicator was developed for use in areas of red parent material, such as Triassic-Jurassic sediments in the Connecticut River Valley, Permian “red beds” in Kansas, clayey red till and associated lacustrine deposits around the Great Lakes, and Jurassic sediments associated with “hogbacks” on the eastern edge of the Rocky Mountains. This indicator also occurs on “Red River” flood plains, such as those along the Chattahoochee, Congaree, Red, and Tennessee Rivers. The most noticeable redox features in red materials are redox depletions and soft manganese masses that are black or dark reddish black (see illustrations on previous pages).



Photo credit: Donald Parizak

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement. Also see Indicator F21 Red Parent Material (page 44). Questions about hydric soil morphologies in red parent materials should be directed to Donald Parizek, Soil Scientist at the Connecticut NRCS office in Tolland.

DEP Particularly Difficult Soils: Indicator TF12: Very Shallow Dark Surface

Technical Description: In depressions and other concave landforms, one of the following:

- If bedrock occurs between 6 in. and 10 in., a layer at least 6 in. thick starting within 4 in. of the soil surface with a value of 3 or less and chroma of 1 or less, and the remaining soil to bedrock must have the same colors as above or any other color that has a chroma of 2 or less.
- If bedrock occurs within 6 in., more than half of the soil thickness must have a value of 3 or less and chroma of 1 or less, and the remaining soil to bedrock must have the same colors as above or any other color that has a chroma of 2 or less.

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator TA6: Mesic Spodic

Technical Description: A layer 2 in. or more thick starting within 6 in. of the mineral soil surface that has a value of 3 or less and chroma of 2 or less and is underlain by either:

- a layer(s) 3 in. or more thick starting within 12 in. of the mineral soil surface that has a value and chroma of 3 or less and shows evidence of spodic development; or
- a layer(s) 2 in. or more thick starting within 12 in. of the mineral soil surface that has a value of 4 or more and chroma of 2 or less and is directly underlain by a layer(s) 3 in. or more thick with a value and chroma of 3 or less that shows evidence of spodic development.

User Notes: This indicator is used to identify wet soils with spodic materials or that meet the definition of a Spodosol in MLRAs 144A and 145 of LRR R and MLRA 149B of LRR S only. The layer that has a value of 4 or more and chroma of 2 or less is typically described as an E or Eg horizon. These typically have color patterns described as stripped or partially stripped matrices. The layer with evidence of spodic development is typically described as a Bh, Bhs, Bhsm, Bsm, or Bs horizon. These layers typically have color patterns or cementation indicative of the accumulation of translocated iron, aluminum, and/or organic matter.




Photo credit: Mark Stolt

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

BBNEP Note: See Appendix C for field criteria to identify spodosols. For additional hydric spodosol morphologies see Indicators S8, S9, IX. A., B., & C. Spodosols.


DEP Particularly Difficult Soils: IX A. Spodosol


Technical Description: Mineral soils having a *spodic horizon* and the following morphology:

A. within 6 inches of the *top of the mineral soil material* have an *E horizon (eluvial horizon)* with 2 percent or more *redoximorphic features* and/or have a *stripped matrix*, that is *directly underlain* by a *spodic horizon* with either:

1. a *Bh, Bhs, or Bs horizon* with 2 percent or more *redoximorphic features*; or
2. a *Bh and/or Bhs horizon* that is *directly underlain*, but within 20 inches of the *top of the mineral soil material*, by a horizon with 2 percent or more *redoximorphic features*;

User Notes: Soils with an *E horizon* that have dark brown to reddish brown colors in the horizon directly underlying it are considered to have evidence of spodic development. The *E horizon* is often discontinuous across the landscape because of natural disturbances. Some soils will have a *dark A or Ap horizon* overlying a dark brown to reddish brown *spodic horizon*. Not all soils that show evidence of spodic development will classify taxonomically as having a *spodic horizon*. These soils are considered problem soils. When soil morphology seems inconsistent with the landscape, vegetation, or observable hydrology, the assistance of an experienced soil or wetland scientist may be needed to determine whether the soil is hydric.





Terms or phrases in *italics* are defined in the Glossary of Terms in Field Indicators for Identifying Hydric Soils in New England, Version 3.

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using these indicators. Additional indicators of hydrology are found in the Regional Supplement.

BBNEP Note: See Appendix C for field criteria to identify spodosols. For additional hydric spodosol morphologies see Indicators S8, S9, TA6, IX B, & C., Spodosols.


DEP Particularly Difficult Soils: IX B. Spodosols

Technical Description: Mineral soils having a *spodic horizon* and the following morphology:

B. beginning within 10 inches of the *top of the mineral soil material* and directly underlying a *dark A* or *Ap horizon* and/or a *shallow E horizon* (or if neither is present, an *O horizon*), there is a *Bh* and/or *Bhs horizon* that is greater than 2 inches thick that is *directly underlain*, but within 20 inches of the *top of the mineral soil material*, by a horizon with 2 percent or more *redoximorphic features*.

User Notes: Soils with an *E horizon* that have dark brown to reddish brown colors in the horizon directly underlying it are considered to have evidence of spodic development. The *E horizon* is often discontinuous across the landscape because of natural disturbances. Some soils will have a *dark A* or *Ap horizon* overlying a dark brown to reddish brown *spodic horizon*. Not all soils that show evidence of spodic development will classify taxonomically as having a *spodic horizon*. These soils are considered problem soils. When soil morphology seems inconsistent with the landscape, vegetation, or observable hydrology, the assistance of an experienced soil or wetland scientist may be needed to determine whether the soil is hydric.

Terms or phrases in italics are defined in the Glossary of Terms in [Field Indicators for Identifying Hydric Soils in New England, Version 3](#).



BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

BBNEP Note: See Appendix C for field criteria to identify spodosols. For additional hydric spodosol morphologies see Indicators S8, S9, TA6, IX. A., & C. Spodosols.

DEP Particularly Difficult Soils: IX C. Spodosols

Technical Description: Mineral soils having a *spodic horizon* and one of the following morphologies:

C. beginning within 10 inches of the *top of the mineral soil material* and directly underlying a *dark A* or *Ap horizon*, is one of the following:



1. an *E horizon* with 2 percent or more *redoximorphic features* and/or a *stripped matrix directly underlain* by a *Bh*, *Bhs*, or *Bs horizon* with 2 percent or more *redoximorphic features*; or
2. a *Bh* and/or *Bhs horizon* that is *directly underlain*, but within 20 inches of the *top of the mineral soil material*, by a horizon with 2 percent or more *redoximorphic features*;

User Notes: Soils with an *E horizon* that have dark brown to reddish brown colors in the horizon directly underlying it are considered to have evidence of spodic development. The *E horizon* is often discontinuous across the landscape because of natural disturbances. Some soils will have a *dark A* or *Ap horizon* overlying a dark brown to reddish brown *spodic horizon*. Not all soils that show evidence of spodic development will classify taxonomically as having a *spodic horizon*. These soils are considered problem soils. When soil morphology seems inconsistent with the landscape, vegetation, or observable hydrology, the assistance of an experienced soil or wetland scientist may be needed to determine whether the soil is hydric.

Terms or phrases in italics are defined in the Glossary of Terms in Field Indicators for Identifying Hydric Soils in New England, Version 3.

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

BBNEP Note: See Appendix C for field criteria to identify spodosols. For additional hydric spodosol morphologies see Indicators S8, S9, TA6, IX. A., & B. Spodosols.



DEP Particularly Difficult Soils: Indicator VIII. Dark Surface Materials

Technical Description: Soils with a matrix chroma of 2 or less that extends to a depth of 20 inches below the top of the mineral soil material, and that have a dark A or Ap horizon (with or without an O horizon) that is directly underlain by a horizon with a matrix value of less than 4, and within 12 inches of the top of the mineral soil material or directly underlying an A or Ap horizon, whichever is shallower, 2 percent or more redoximorphic features that extend to:

- A. a depth of 20 inches below the top of the mineral soil material; or
- B. a depleted or gleyed matrix, whichever is shallower;





Photo Credit: Jim Turrene

BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

VIII. A. and B. Dark Mineral Soils




Key for Soil Textures

 Slightly to partially decomposed organic matter	 Mucky mineral soil	 Any mineral soil texture
 Well decomposed organic matter	 Loamy fine sand or coarser soil textures	

DEP Particularly Difficult Soils: Indicator XI. A. Redox Depletions (Any Texture)

Technical Description: Soils that do not have a *spodic horizon* and beginning within 10 inches of the *top of the mineral soil material* and directly underlying a *dark A* or *Ap horizon* (with or without an *O horizon*) is a horizon with 5 percent or more *redox depletions*, and within 20 inches of the *top of the mineral soil material* there is a horizon with a *depleted* or *gleyed matrix* (for soils with *moderate to strong structure*, the *matrix color* and percent *redox depletions* are recorded for *ped interiors*).

Terms or phrases in italics are defined in the Glossary of Terms in [Field Indicators for Identifying Hydric Soils in New England, Version 3](#).




BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

DEP Particularly Difficult Soils: Indicator XII Chroma 3 with Very Dark Surface (Any Texture)

Technical Description: Soils that have a *very dark A* or *Ap* horizon less than 10 inches thick (with or without an *O* horizon) that are *directly underlain* by a horizon with a *matrix color* due to wetness of chroma 3 or less, with 10 percent or more *redoximorphic features*; and within 6 inches of the *top of the mineral soil material* have 2 percent or more *redoximorphic features*; and within 18 inches of the *top of the mineral soil material* have 2 percent or more *redox depletions*.


Terms or phrases in italics are defined in the Glossary of Terms of Field Indicators for Identifying Hydric Soils in New England, Version 3.



BBNEP Note: Look for a second confirmation of wetland hydrology, as listed on the DEP Form, when using this indicator. Additional indicators of hydrology are found in the Regional Supplement.

Appendix A Major Land Resource Areas (MLRA) and Land Resource Regions (LLR) for DEP Particularly Difficult Soils

Massachusetts is predominately in LLR R. Region and subregion boundaries are depicted in the figure below as sharp lines. However, climatic conditions and the physical and biological characteristics of landscapes do not change abruptly at the boundaries. In reality, regions and subregions often grade into one another in broad transition zones that may be tens or hundreds of miles wide. The lists of wetland indicators presented in the Regional Supplement for the Northcentral and Northeast region, as depicted in this guide, may differ between adjoining regions or subregions. In transitional areas, the investigator must use experience and good judgment to select the supplement and indicators that are appropriate to the site based on its physical and biological characteristics. Wetland boundaries are not likely to differ between subregions in transitional areas, but one subregion criteria may provide more detailed treatment of certain problem situations encountered on the site. If in doubt about which criteria to use in a transitional area, apply all available indicator criteria and compare the results.



Location of MLRAs 144A and 145 in LRR R and MLRA 149B in LRR S.

BBNEP Note: Users in the Buzzards Bay watershed are in the transition zone of MLRA 144A and MLRA 149B. use indicators for both regions in this area.

Appendix B: Thickness Criteria for DEP Particularly Difficult Soils

It is permissible to combine certain hydric soil indicators if all requirements of the individual indicators are met except thickness (see Hydric Soil Technical Note 4, http://soils.usda.gov/use/hydric/ntchs/tech_notes/index.html). The most restrictive requirements for thickness of layers in any indicators used must be met. Not all indicators are possible candidates for combination. For example, indicator F2 (Loamy Gleyed Matrix) has no thickness requirement, so a site would either meet the requirements of this indicator or it would not. Table B1 lists the indicators that are the most likely candidates for combining in the region.

Table B1: Minimum thickness requirements for commonly combined indicators in the Northcentral and Northeast Region

Indicator	Thickness Requirement
S5 – Sandy Redox	4 in. thick starting within 6 in. of the soil surface
S7 – Dark Surface	4 in. thick starting within 6 in. of the soil surface
F1 – Loamy Mucky Mineral	4 in. thick starting within 6 in. of the soil surface
F3 – Depleted Matrix	6 in. thick starting within 10 in. of the soil surface
F6 – Redox Dark Surface	4 in. thick entirely within the upper 12 in.
F7 – Depleted Dark Surface	4 in. thick entirely within the upper 12 in.

Table B2 presents an example of a soil in which a combination of layers meets the requirements for indicators F6 (Redox Dark Surface) and F3 (Depleted Matrix). The second layer meets the morphological characteristics of F6 and the third layer meets the morphological characteristics of F3, but neither meets the thickness requirement for its respective indicator. However, the combined thickness of the second and third layers meets the more restrictive conditions of thickness for F3 (i.e., 6 in. starting within 10 in. of the soil surface). Therefore, the soil is considered to be hydric based on the combination of indicators.

Table B2. Example of a soil that is hydric based on a combination of indicators F6 and F3.

Depth (inches)	Matrix Color	Redox Concentrations			Texture
		Color	Abundance	Contrast	
0 – 3	10YR 2/1	--	--	--	Loamy/clayey
3 – 6	10YR 3/1	7.5YR 5/6	3 percent	Prominent	Loamy/clayey
6 – 10	10YR 5/2	7.5YR 5/6	5 percent	Prominent	Loamy/clayey
10 – 14	2.5Y 4/2	--	--	--	Loamy/clayey

Appendix B: Thickness Criteria for DEP Particularly Difficult Soils, cont.

Another common situation in which it is appropriate to combine the characteristics of hydric soil indicators is when stratified textures of sandy (i.e., loamy fine sand and coarser) and loamy (i.e., loamy very fine sand and finer) material occur in the upper 12 in. of the soil. For example, the soil shown in Table B3 is hydric based on a combination of indicators F6 (Redox Dark Surface) and S5 (Sandy Redox). This soil meets the morphological characteristics of F6 in the first layer and S5 in the second layer, but neither layer by itself meets the thickness requirement for its respective indicator. However, the combined thickness of the two layers (6 in.) meets the more restrictive thickness requirement of either indicator (4 in.).

Table B3. Example of a soil that is hydric based on a combination of indicators F6 and S5.

Depth (inches)	Matrix Color	Redox Concentrations			Texture
		Color	Abundance	Contrast	
0 – 3	10YR 3/1	10YR 5/6	3 percent	Prominent	Loamy/clayey
3 – 6	10YR 4/1	10YR 5/6	3 percent	Prominent	Sandy
6 – 16	10YR 4/1	--	--	--	Loamy/clayey

Appendix C: Spodosol Field Criteria

When an Ap is present look for the following colors directly below the Ap (If an E-horizon is present look for these directly below the E):

- a. a hue of 5YR or redder (i.e. 2.5YR, etc); or
- b. a hue of 7.5YR, color value 5 or less and chroma 4 or less (a hue of 7.5YR with value 5 and chroma 6 does not qualify without additional chemical properties); or
- c. a hue of 10YR or neutral and color value and chroma 2 or less; or
- d. a color of 10YR 3/1.

Bs – value and chroma are more than 3

Bhs - both the value and chroma are 3 or less

Spodosols are a problem soil. Check DEP Difficult Soils: Evergreen Forest Soils, and DEP Particularly Difficult Soils S8, S9, and TA6.



Photo credit: John Rockwell. This is an example of a spodosol altered by agricultural activity. This activity has obliterated the E horizon and the top of the Bhs horizon. This soil meets the criteria for DEP Particularly Difficult Soils Indicator TA6.

BBNEP Note: The identification of spodosols can be difficult. If the site has been subject to some kind of agricultural activity, the E-horizon and spodic horizon may have been mixed with the A horizon to form an Ap horizon, sometimes known as a plow layer. Look below the Ap to see if the soil has a reddish tinge.

Appendix D: Mucky Mineral Field Identification for DEP Particularly Difficult Soils - Determining the texture of soil materials high in organic carbon.

Material high in organic carbon could fall into three categories: organic, mucky mineral, or mineral. In lieu of laboratory data, the following estimation method can be used for soil material that is wet or nearly saturated with water. This method may be inconclusive with loamy or clayey textured mineral soils. Gently rub the wet soil material between forefinger and thumb. If upon the first or second rub the material feels gritty, it is mineral soil material. If after the second rub the material feels greasy, it is either mucky mineral or organic soil material. Gently rub the material two or three more times. If after these additional rubs it feels gritty or plastic, it is mucky mineral soil material; if it still feels greasy, it is organic soil material.



Photo credit: <http://www.ohiowineandmore.com/>

Appendix E: Redox Features- Faint vs. Distinct

Compare the matrix color to the redox feature color. The contrast is distinct if: (Note: Regardless of the magnitude of hue difference, where both colors have value ≤ 3 and chroma ≤ 2 , the contrast is faint.)

Threshold for Distinct		
Δ Hue	Δ Value	Δ Chroma
0	≤ 2	>1 to <4
0	>2 to <4	<4
1	≤ 1	>1 to <3
1	>1 to <3	<3
2	0	>0 to <2
2	>0 to <2	<2

Conversely, faint is evident only on close examination. The contrast is faint if:

Upper Threshold for Faint		
Δ Hue	Δ Value	Δ Chroma
0	≤ 2	≤ 1
1	≤ 1	≤ 1
2	0	0
Hue	Value	Chroma
Any	≤ 3	≤ 2

Any feature above the upper threshold for faint features would be considered either distinct or prominent. If an indicator requires distinct or prominent features then those features at or below the faint threshold do not count.

Appendix F: ACOE Memo regarding use of "Field Indicators for Identifying Hydric Soils in New England, Version 3



REPLY TO
ATTENTION OF

DEPARTMENT OF THE ARMY
NEW ENGLAND DISTRICT, CORPS OF ENGINEERS
699 VIRGINIA ROAD
CONCORD, MASSACHUSETTS 01742-2751

CENAE-R-PT

March 11, 2011

MEMORANDUM FOR THE RECORD

SUBJECT: *Field Indicators for Identifying Hydric Soils in New England (Version 3)*. New England Interstate Water Pollution Control Commission (NEIWPCC), April 2004

1. This guide is an update of the previous version released in 1998. It clarifies and refines the 1998 version based on extensive field testing. This guide is currently the best available reference of its kind in New England and is specifically developed for New England soils.
2. This version of this guide is widely used by state and Federal agency staff and the consulting community. It is a standard reference for regulatory programs throughout much of New England.
3. This field guide provides an important resource to use in problem and disturbed situations where Chapter 5 "Difficult Wetland Situations in the Northcentral and Northeast Region" of the current *Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region* is applicable. When properly used, this field guide provides results that help make determinations in these areas.
4. The Environmental Resource Section staff of the Policy Analysis and Technical Support Branch in the Regulatory Division use this field guide and continue to encourage its use by Corps staff and other wetland practitioners for Chapter 5 circumstances.

A handwritten signature in black ink, appearing to read "Ruth M. Ladd".

RUTH M. LADD
Chief, Policy Analysis and
Technical Support Branch

Appendix G – Observing and Recording Hue, Value and Chroma

All colors noted in this supplement refer to moist Munsell® colors (Gretag/Macbeth 2000). Do not attempt to determine colors while wearing sunglasses or tinted lenses. Colors must be determined under natural light and not under artificial light.

Chroma

Soil colors specified in the indicators do not have decimal points (except for indicator A12); however, intermediate colors do occur between Munsell chips. Soil color should not be rounded to qualify as meeting an indicator. For example, a soil matrix with a chroma between 2 and 3 should be recorded as having a chroma of 2+. This soil material does not have a chroma of 2 and would not meet any indicator that requires a chroma of 2 or less.

Hue and Value

Hue and value should be rounded to the nearest color chip when using the indicators. For example, if the color is in between value of 3 and 4 it should be rounded and not excluded from meeting either F3 Depleted Matrix or F6 Redox Dark Surface because it is in between values. If the value is closer to a 3 then F6 or some other dark surface indicator should be considered and if it is closer to 4 then F3 or some other depleted matrix indicator should be considered.

Timing - not too wet, and not too dry

Always examine soil matrix colors in the field immediately after sampling. Ferrous iron, if present, can oxidize rapidly and create colors of higher chroma or redder hue. In soils that are saturated at the time of sampling, redox concentrations may be absent or difficult to see, particularly in darkcolored soils. It may be necessary to let the soil dry to a moist state (5 to 30 minutes or more) for the iron or manganese to oxidize and redox features to become visible.

Postscript: Indicators of Hydrology

Primary Indicators (minimum of one required)

Surface Water	Water-Stained Leaves
High Water Table	Aquatic Fauna
Saturation	Oxidized Rhizospheres on Living Roots
Hydrogen Sulfide Odor	Sediment Deposits
Water Marks	Drift Deposits
Presence of Reduced Iron	Algal Mat or Crust
Recent Iron Reduction in Tilled Soils	Iron Deposits
Thin Muck Surface	Inundation Visible on Aerial Imagery
Sparsely Vegetated Concave Surface	

Secondary Indicators (minimum of two required)

Surface Soil Cracks	Drainage Patterns
Moss Trim Lines	Dry-Season Water Table
Crayfish Burrows	Saturation Visible on Aerial Imagery
Stunted or Stressed Plants	Geomorphic Position
Shallow Aquitard	Microtopographic Relief
FAC-Neutral Test	

BBNEP Note: For details on these hydrology indicators, see the DEP Delineation Manual and the Regional Supplement.



Buzzards Bay National Estuary Program
2870 Cranberry Highway
East Wareham, MA 02538
#508-291-3625 x 14
www.buzzardsbay.org